



Virginia Commonwealth University  
**VCU Scholars Compass**

---

Biology and Medicine Through Mathematics  
Conference

---

2017

May 18th, 12:30 PM - 1:00 PM

# Dynamics of Fluorescent Imaging in Glob-Driven Breakup

Lan Zhong

*University of Delaware*, [lanzhong@udel.edu](mailto:lanzhong@udel.edu)

Christiaan Ketelaar

*University of Delaware*, [ketelaar@udel.edu](mailto:ketelaar@udel.edu)

Richard J. Braun

*University of Delaware*, [rjbraun@udel.edu](mailto:rjbraun@udel.edu)

*See next page for additional authors*

Follow this and additional works at: <http://scholarscompass.vcu.edu/bamm>



Part of the [Life Sciences Commons](#), [Medicine and Health Sciences Commons](#), and the [Other Applied Mathematics Commons](#)

---

<http://scholarscompass.vcu.edu/bamm/2017/thursday/2>

This Event is brought to you for free and open access by the Dept. of Mathematics and Applied Mathematics at VCU Scholars Compass. It has been accepted for inclusion in Biology and Medicine Through Mathematics Conference by an authorized administrator of VCU Scholars Compass. For more information, please contact [libcompass@vcu.edu](mailto:libcompass@vcu.edu).

---

**Presenter Information**

Lan Zhong, Christiaan Ketelaar, Richard J. Braun, Carolyn G. Begley, and Peter Ewen King-Smith

# Dynamics of Fluorescent Imaging in Glob-Driven Breakup

L. Zhong, C. F. Ketelaar, R.J. Braun, P.E. King-Smith, C.G. Begley

February 28, 2017

The tear film is the very thin film ( $3\mu\text{m}$ ) left behind after a blink. The dynamics of tear film breakup (TBU) is important for understanding Dry Eye Syndrome (DES), which affects millions, and can help clarify the mechanisms of DES for ophthalmologists. Recently, a different type of breakup is observed, where tear film breakup (TBU) happens under or around a thick circular area of lipid in a second after blink; the thick lipid corresponds to a glob. Evaporation alone is too slow to offer a complete explanation of this type of breakup. Instead, a divergent flow driven by Marangoni effect is proposed to explain the rapid thinning. We examined the Marangoni-driven TBU hypothesis in a one dimensional and an axisymmetric models by assuming the thick lipid (glob) has higher surfactant concentration, thus a lower surface tension. In both models, the thick lipid spreads out immediately due to Marangoni effect, subsequently, a strong outwards flow dragged by the corresponding shear stress leads to TBU within a second. In addition, transport of osmolarity and fluorescein are also studied in the models. TBU happens while a weak osmotic flow flows across the ocular surface. The distribution of salt ions and fluorescein can be affected by the evaporation but are mostly determined by the Marangoni flow. The models predict that smaller globs or thinner film will decrease TBU time (TBUT). TBU will be seen under the glob, when glob is small, otherwise TBU occurs outside the glob. This study will help provide insights about mechanism of rapid thinning and how the properties of the glob will affect TBUT and visualization of TBU.